

Type III

MAR 23 1959

Rectal Swabs

Tubes 1371 as attached sheets

Procedure 0.4 mls of fluid added to each of 3 tubes.

<u>Date of Rectal Swab</u>	<u>Tubes</u>	<u>3/25</u>	<u>3/26</u>	<u>3/27</u>	<u>3/29</u>	<u>3/30</u>	<u>3/31</u>
	<u>B.</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>6</u>	<u>7</u>	<u>8</u>
2/10/59 free	190	0	0	0	0	0	0
	191	0	0	0	0	0	0
	192	0	0	0	0	0	0
2/12/59 2 days	193	0	0	0	0	0	0
	194	0	0	0	0	0	0
	195	0	0	0	0	0	0
2/14/59 4 days	196	0	0	0	0	0	0
	197	0	0	0	0	0	0
	198	0	0	0	0	0	0
2/16/59 6 days	199	1-3	4 off	} Pooled Sewed			
	200	2-3	4 off				
	201	1-2	4 off				
2/19/59 9 days	202	0	2-3	4 off	} Pooled Sewed		
	203	few	3-4	4 off			
	204	0	1-2	4 off			
2/23/59 13 days	205	0	1-2	3-4 off			
	206	0	0	0	0	0	0
	207	2-3	3-4 off	Sewd.			
3/3/59 21 days	208	0	0	0	0	0	0
	209	0	0	0	0	0	0
	210	0	0	0	0	?	?
						} Sewed to loss	
						210	0 0

Passage 4/6/59
 4/7 4/8 4/9/59
 1 2 3 5

Type III

MAR 23 1959

Throat Swabs.

Tubes. 1371 as overleaf

Procedure 0.4 ml of fluid added to each of 3 tubes.

<u>Date of Throat Swab</u>	<u>Tubes</u>	<u>3/25</u>	<u>3/26</u>	<u>3/27</u>	<u>3/29</u>	<u>3/30</u>	<u>3/31</u>
	B.	2	3	4	6	7	8
2/10/59	211	0	0	0	0	0	0
he	212	0	0	0	0	0	0
	213	0	0	0	0	0	0
2/12/59	214	0	0	0	0	0	0
2 days	215	0	0	0	0	0	0
	216	0	0	0	0	0	0
2/14/59	217	0	0	0	0	0	0
4 days	218	0	0	0	0	0	0
	219	0	0	0	0	0	0
2/16/59	220	0	0	0	0	?	0
6 days	221	0	0	0	0	0	0
	222	0	0	0	0	0	0
2/19/59	223	0	0	0	0	0	0
9 days	224	0	0	0	0	0	0
	225	0	0	0	0	0	0
2/23/59	226	0	0	0	0	0	0
13 days	227	0	0	0	0	0	0
	228	0	0	0	0	0	0
	229	0	0	0	0	0	0
3/3/59	229	0	0	0	0	0	0
21 days	230	0	0	0	0	0	0
	231	0	0	0	0	0	0

larvae 4/6/59
 4/7 4/8 4/9 4/11
 ↓ 2 3 5

→ Served to pos
 → 220 0 0 0

type III Group.

APR 13 1959

Material Original fluid of R.S. of 2/16/59

Serum Polio III antiserum R63 r64 1:10

tubes 1377 of 4/7/59

		4/15 2	4/16 3	4/17 4	4/18 5	4/20 7	
Fluid + equal vol serum Polio III	111	0	0	0-1	2-3 vac. 40%	7	Save
	112	0	0	few	0-1	2-40%	
	113	0	0	0	0	containing some Polio III	
+ equal vol diluent	114	2-3	40%				+ ?
	115	2-3	40%				
	116	2-3	40%				
10 ⁻¹	117	0-1	3-40%				10 ^{3.0} /0.2 ml
	118	1-2	3-40%				10 ^{5.7} /ml.
10 ⁻²	119	0	2-3	40%			
	120	0	1-2	40%			
10 ⁻³	121	0	0	0	0	0	0
	122	0	0	2-3	40%		
10 ⁻⁴	123	0	0	0	0	0	0
	124	0	0	0	0	0	0

Sed type III

APR 28 1959

To identify the virus in the Rectal Swabs of 2/19/59 & 2/23/59

Material MKP of R.S of 2/19 and 2/23.

Seum Polio III antiserum R.63 + R.64 of 4/1/55 1:10

Tubes T 379 of 4/21/59.

	tubes	4/30	5/1	5/2	5/4	5/6	
	A.	2	3	4	6	8	
<u>2/19/59 R.S. MKP 10⁻⁴ + Polio III</u>	109	0	0	0	0	0	} Keep
	110	0	0	0 _{var}	0	0	
	111	0	0	0	0	0	
	+ diluent		112	3	40%		
	113	3-4	40%				
	114	3	40%				
<u>2/23/59 R.S. MKP 10⁻⁴ + Polio III</u>	115	0	0	0	0	0	} Keep
	116	0	0	0	0	0	
	117	0	0	0 _{var}	0 _{var}	0	
	+ diluent		118	3-4	40%		
	119	3-4	40%				
	120	3-4	40%				